

THE PARA-TYMPANIC ORGAN (PTO) of VITALI : a documental * continuing forgotten request to the SCIENCE

THE PARA-TYMPANIC ORGAN (PTO) of VITALI : a documental *
continuing forgotten request to the SCIENCE

** Supposed PTO function's "evidence based " records – twenty
years woodpigeons' Europe migration monitoring – from the
field , updated 2018*

Enrico Cavina

*Club Italiano del Colombaccio (member SEEN / Bird Migration
Research Foundation)*

ABSTRACT

*The writing architecture of the present paper is built for on-
line reading .*



The provocative image is exhaustive of the Abstract's text

A cry of indignation to the Science

***Key-words : PT0 – wood pigeon's migration – counts – anatomy-
waves & peaks – Air Pressure (hPa) – isobars- abiotic factors
– “evidence based” data – Research***

***FOR ANY PROBLEM ON THE WEB-LINKS , please, COPY and PASTE on
the window of GOOGLE search motor .***

Italian anatomist, Vitali was professor of Anatomy at Pisa. Vitali described the constant presence of small ganglia on orbital branches of the sphenotemporal ganglion in humans. He reported the discovery of the paratympanic organ, a sense organ in the middle ear of birds. He was the first to postulate that the paratympanic and spiracular sense organs were homologous organs that originate from homologous placodes.

Giovanni VITALI
1836 Portico della Chiesa AR (I) 1903 Pisa (I)



INTRODUCTION

Unfortunately the interest in the specific research on the sensitive organ Paratympanic of Vitali (PTO) (1-2-3-4-5-6) , remains as “zero” despite many typologies of open research’s fields over PTO and related questions and still unsolved answers from the Science .

It is basic to say “ *PTO is used daily by enormous number of living animals (200-300 billions of birds) but represents currently one of the major unsolved mysteries and challenges in sensory physiology of vertebrates* ” (8) (Giannessi F. and co-Aut.,2013).

We have explored many items concerning PTO by retroactive and

real-time analysis of “evidence based” data from Bibliography, Web-bibliography, Archives and from the field as previously published (35-35-41) .

The basic question is as following : “ *why 200-300.000 woodpigeons in stop-over started all together in few minutes by a mass-takeoff , continuing transcontinental migration ?* ” as happened 31 October 2017 from Mesola forest (Ferrara-Italy) fully documented and by the video <https://youtu.be/pGK6z9SY8Cg> (Denis Bianchi -2017) and our forecast 4 days before the event.

Mass-continuing-fly at same altitude (altimetric function) and isobaric corridors as by video 2018 from Pamplona (Spain) <https://youtu.be/4Tb5kGRagtg> –<https://youtu.be/ty6pdmkSdGo>

What, where is the finger pushing the bottom of the takeoff , and by what typology of command ? It happened also in other sites and past years in Italy and Europe . Many elements suggest that the focus must be on PT0 and its sensitive barometric altimetric function , as also we supposed by retroactive archives-research on woodpigeons (and other birds) autumn migration (41).

The text of our precedent short-paper “ *PT0 in migratory birds: research’s failure or challenge ?* ” (<http://www.scienceheresy.com/ornithologyheresy/Cavina2016.pdf> 2016 –) (38) seems still valid but -by our personal opinion- the balance is lowering to permanent failure despite so much fascinating topic : the text of mentioned short-paper should be considered as introductive to the present updating and discussion .

MATERIALS and METHODS

RESULTS

The present -2018- collected data integrate the precedent data collected in 2014 (<http://www.scienceheresy.com/ornithologyheresy/Cavina2015.pdf> and <https://www.researchgate.net/publication/328336987> 2018 -) (41)

The supposed barometric function of PT0 in wild migratory birds is strictly linked with the decision-making-time to the migration mass take-off and/or birds' populations departing in concentrated waves from origin and stop-over sites

Many data (biotic and abiotic) factors concerning this item have been reported and discussed in a first paper on-line (*"Decision making of autumn migrations of woodpigeons (Columba palumbus) in Europe: analysis of the abiotic factors and atmospheric pressure changes"* – November 2014 - <http://www.scienceheresy.com/ornithologyheresy/Cavina2015.pdf>) (41) .

All the data-results also in graphics of analysis on many abiotic factors and specifically on the changes in Air Pressure values controlled along the migration flyways , seem –also at present time November 2018 in updating review – as clearly indicative concerning the relationship between migration takeoff's decision-making and jumps of Air Pressure before the take-off.

<https://plus.google.com/photos/103942035281038458760/albums/6086432829554328737>

Many of details reported and discussed are connected with data – updated 2018- collected in Falsterbo (SE) and Pyrenees (FR) and authorized to be used by direct Official local Institutions (personal communication emailed 2014).

These data concerning migration waves(**W-3 days more**) and peaks (**PK-1-2 days**) –updated November 2018 – are here reported as following and are in relation

- with the jumps (**more 10 hPa**) of Air Pressure (AP)
- and 12-24 h before takeoff (**j-1**) or 24-48 h before takeoff (**j-2**) – none jump (**j-0**)
- and total/year as 1999 (148.360)

All the meteo-data as basic of the retroactive analysis ,available to be documentative , are on-line by many Meteo Websites , but in our work mostly by

“Weather History – wunderground “

<https://www.wunderground.com/history/>

“ www.wetterzentrale.de “ as by the Link and isobaric maps

<http://www.wetterzentrale.de/en/reanalysis>.

—
By these Links it's possible to check all the old or ancient meteo-data day by day, hour by hour, locality by locality, year by year .

—
—
—
Why the focus on the Air Pressure in atmosphere ? The answer is : high values of Air Pressure are warranty for stable isobaric altitudinal corridors in migration's flyway without turbulences and connected economy of energies .

The focus is on “PEAKS” & “waves” on the following analysis .
(41)

*** *** ***

SWEDEN south (13.660.418 woodpigeons counted in 20 years)

Falsterbo Bird Observatory – data from ARCHIVES –

https://www.falsterbofagelstation.se/index_e.html

▪ **1999** (148.360)

TWO waves – 6-7-8 Oct. (W-34643) (j-1) – 12-13-14 Oct.
(W-43.835) (j-0)

▪ **2000** (333.125)

TWO waves-ONE peak – 15-16-17 Oct. (W-40540) (j-1) - 27-28
Oct. (PK-54300) (j-1) – 2-3-4 Nov. (W-79445) (j-1)

▪ **2001** (208.815)

ONE peak –THREE waves – 6-7 Oct. (PK -37600) (j-1) -11-12-13
Oct. (W-32240) (j-2) – 17-18-19 Oct. (W – 79445) (j-1)

▪ **2002** (263.351)

THREE waves –ONE peak – 6-7-8-Oct. (W-46800) (j-1) -10-11-12
Oct. (W -44400) (j-2) -

20-21 Oct. (PK -60900) (j-1) - 29-30-31 Oct. (W-40300) (j-1)

▪ **2003** (144.496)

ONE wave – 12-13-14 Oct. (W-69800) (j-1)

▪ **2004** (249.560)

THREE peaks – ONE wave – 9-10-11 Oct. (W-36350) (j-1) - 19-20
Oct. (PK -38760) (j-1) -

27 Oct. (PK-43950) (j-1) - 2 Nov. (PK-56800) (j-2)

▪ **2005** (421.710)

TWO waves – ONE peak – 11-12-13 Oct. (W-106000) (j-1) - 15-16

Oct.(PK 165900) (j-2)- 23-24-25.26 Oct. (W – 69135)(j-1)

▪ **2006** (429.560)

ONE wave – TWO peaks – 13-14-15-16 Oct. (W- 218.416) – 25-26 Oct. (PK-84800)(j-1)- 2 Nov. (PK- 31100)(j-2)

▪ **2007** (254.286)

ONE wave – TWO peaks – 4-5-6- Oct (W -47130)(j-0) – 13 Oct.(PK – 71360)(j-1)-19-20 Oct. (PK-62070)(j-2)

▪ **2008** (407.702)

THREE peaks – ONE wave – 7-9 Oct.(PK- 78030)(j-1) - 15-16-17-18 Oct. (W-138535)(j-2)- 24-25 Oct.(PK -70040)(j-2) – 29-30 Oct.(PK-55650) (J-1)

▪ **2009** (314.615)

ONE peak – TWO waves – 9-10 Oct. (PK – 61150) (j-1) – 13-14-15 Oct. (W-120650)(j-0)- 27-28-29 Oct-(W- 44800)(J-2)

▪ **2010** (258.582)

THREE waves – 28-29-30 Sept.-1 Oct. (W-67500) (j-2) – 9-10-11 Oct.(W-79700) (J-1)-

15-16-17-18 Oct. (W-59600) (j-1)

▪ **2011** (355.655)

THREE peaks – 8-9 Oct- (PK-156300) (j-1) – 13-14 Oct. (PK-105500)(j-1) – 21 Oct. (PK- 25600)(j-2)

▪ **2012** (491.833)

TWO waves – 10-11-12 Oct. (W- 204600)(j-1)- 19-20-21 Oct. (W- 198990) (j-1)

▪ **2013** (844.710)

THREE peaks – TWO waves – 11-12 Oct. (PK-450800 ***) (j-1) –

15-16-17-18 Oct. (W-103990) (j-2) – 25 Oct. (PK-43200)(j-1)
– 31 Oct. (PK -30500)(j-2) – 5-6-7 Nov. (W-95000) (j-2)

▪ **2014** (562.360)

TWO waves – ONE peak – 11-12-13-14 Oct. (W-111600)(j-1) –
21-22-23 Oct.(W-162900)(j-2) – 28 -30 Oct. (PK -110500) (j-1)

▪ **2015** (684.087)

THREE peaks – 10-11 Oct. (PK-332500)(j-1) – 19-20 Oct.
(PK-88100)((j-1) – 26-27 Oct. (PK-77700) (j-1)

▪ **2016** (779.920)

THREE waves – 3-4-5-6-7-8 Oct. (W-321100)(j-1) – 12-13-14 Oct.
(W-131110)(j-1)-



20-26 Oct (7gg) (W-262.870) (j-2)



▪ **2017** (494.300)

TWO waves – ONE peak – 6-7-8-9 Oct. (W-87400)(j-1)- 23-24 Oct
(PK-212500)(j-1)-27-28-29-30 Oct. (W-87200)(j-1)

▪ **2018** (379.610)

TWO waves – 4-7 Oct. (W-44600)(j-1) – 11-16 Oct.(6days)
(W-218900) (j-2)





Comment

In Falsterbo (SE) from 1999 to 2018 (20 years) the Observers counted Woodpigeons in transit

- 660.418 (average/y 683.020 – min.144.496 – max.844.710)
- 134.943 during 60 waves/peaks (average 102.249) – 45{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of 13.660.418
- 529.524 during 34 waves (days more than 2 – average 103.809) 25{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of 13.660.418 and 57{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of 6.134.943 (total of waves and peaks)

- 605.410 during 26 peaks (1-2 days – average 100.208) –
19{a6be835d09a84c463fa955f66368ad74671de57ffb293aa268345
95a88d1abe5} of 13.660.418 and
42{a6be835d09a84c463fa955f66368ad74671de57ffb293aa268345
95a88d1abe5} of 6.134.943 (total of waves and peaks)

Comparing the average between numbers of Woodpigeons recorded during Waves (103.809) and Peaks (100.208) the difference is poor.

During first 10 years (1999-2008) Woodpigeons counted 2.860.965 (average/y 286.096)

During second 10 y (2009-2018) Woodpigeons counted 5.165.672 (average/y 516.567)

during the second ten years (2009-2018) the increment has been 78,8

{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5}

The PC sistema can have changed some numbers during “insert” : some mistakes are possible

GRAPHICS (2014) **at**
<https://plus.google.com/photos/103942035281038458760/albums/6086432829554328737>

Concerning the specific focus on Air Pressure changes as stimulus to depart (decision making to takeoff by peaks/waves) for Migration (timing from origin and stop-over

sites) –apart all the data related to all abiotic and biotic factors as many reported in graphics- we can underline some conclusions previously reported (...Cavina) and can be assimilated to the current updating (2014-2018).

Air Pressure (AP) : changes in **global growth** (1° - 30°) in 92.62{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of the peaks/waves , variations in growth of over 10 hPa difference in 72,72{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} (48-24 h prior to take-off) and 78 , 72{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} in the 18 hours before, to be regarded as the dawn in the study area occurs at around 7 am and the woodpigeons typically move from the woods about 30'-60 'after sunrise. About the time of the take-off and other environmental factors play important roles and temporal scope of an overnight stop, such as night-time temperature, humidity, clear skies or cloudy, rain, wind, moon phase.□The values of their base at 7-11 pm (Twilight Night) the night before the take-off point to a Atmospheric pressure in hPa between 1011-1040 53.43{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of peak/waves .□The same values in the morning of the take-off and transit (7-11 am) show that 75.45{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of the peaks/waves with differential values higher than 1010 hPa.

Concerning other **abiotic factors** and their influence on the migration-decision-making (**Falsterbo**) other data (1999-2013) are available .

1) Weather conditions:
74.46{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of the peaks/waves correspond to slightly cloudy or clear days.□

2) Wind – strength – 41 of 47 peaks/waves with winds of force

between 6-25 km / h, then mostly moderate and evenly distributed properly in the days of transit in peaks/waves.□

3) Wind – direction –
42.55{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of the peaks/waves with favorable winds predominantly from the North and East on the day of the passage;

38.29{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of the peaks with unfavorable winds predominantly from the south and west;

14.89{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of the peaks under conditions of moderate variability and / or calm winds.

□4) Visibility: 6 - 10 km in
'89.36{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of peaks/waves.□

5) Length of Day: calculated in mid-October between 10 h and 10h 45 '.

□6) Temperature:-average temperature between 6-20 ° on average, mild- 89,
36{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of peaks□/waves

7) Humidity: prevails in absolute terms between 60 -100{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5}.□

8) Moon: percentage of the lunar surface illuminated over 40{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} in
61,70{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of peaks/waves (statistical significative data)

FRANCE south

PYRENEES – by GIFS –data from ARCHIVES

<https://www.palombe.com/migration/comptages.php>

1999-2018 : woodpigeons 31.169.768 (*average/y 1.558.488 – min.1.028.000 – max. 2.820.263*) passed through Pyrenees mountains in last 20 years

▪ **1999** (2.787.626)

THREE peaks -16 Oct.(PK-108197)(j-1) – 26-27 Oct.(PK-1907721)(j-1)- 4 Nov.(PK -358794)(j-1)

▪ **2000** (556.987)

TWO waves – ONE peak – 18-19-20 Oct. (W-434891)(j-1)- 23-24-25 Oct.(W-689783((j-2)- 27-28 Oct.(PK-381373)(j-0)

▪ **2001** (828.783)

TWO peaks –ONE wave – 16 Oct.(PK-223442)(j-1) – 24-25-26-27 Oct.(W-675391)(j-1)- 2-3 Nov.(W-300870)(j-2)

▪ **2002** (2.052461)

TWO peaks –ONE wave – 19-20 Oct.(PK-237723)(j-2) – 27-28-29 Oct.(W-759472)(j-1) – 5-6 Nov. (PK-485769)(j-1)

▪ **2003** (1.624.152)

THREE peaks – 22 Oct.(PK-56050)(j-2)- 25 Oct.(PK-274100)(J-2) -
3 Nov.(PK-1009490)(j-1)

▪ **2004** (1.820.180)

THREE peaks – 22-23 Oct.(PK-353168)(j-1) – 30-31
Oct.(PK584070)(j-1)- 3 Nov. (PK-139692)(j-2)

▪ **2005** (2.477.468)

ONE peak-TWO waves – 17 Oct.(PK-114122)(j-2)- 24-25-26-27
Oct.(W-1037592)(j-1)- 30-31 Oct.-1-2-3 Nov.(W-1101595)(j-2)

▪ **2006** (1.520.491)

TWO waves – 25-26-27-28 Oct.(W-799537)(j-1)- 2-3-4
Nov.(W-563868)(j-2)

▪ **2007** (1.643.274)

TWO waves – ONE peak – 19-20-21-22 Oct.(W-657146)(j-2)-27-28
Oct.(PK-416226)(j-2)-6-7-8 Nov.(W-205195)(j-1)

▪ **2008** (2.208.369)

TWO waves ONE peak – 17-18-19 Oct.(W-1439480)(j-1) –
23-24-25-26 Oct.(W- 727626)(j-2)- 4 Nov. (PK-205195)(j-0)

▪ **2009** (1.310.538)

ONE peak ONE wave – 18-19 Oct.(PK 88479)(j-2) – 26-27-28-29-30
Oct.(W-1108525)(j-1)

▪ **2010** (1.028.000)

TWO peaks – 21-22 Oct.(PK-508539)(j-1) – 26-27
Oct.(PK-307565)(j-1)

▪ **2011** (1.440.469)

TWO peaks ONE wave – 21-22 Oct.(PK-315616)(j-1)- 31 Oct.(PK
606878)(j-1)-

9-10-11 Nov. (W-390316) (j-1)

▪ **2012** (2.449.624)

TW0 peaks ONE wave – 18 Oct. (PK-158069) (j-1) - 28-29-30-31 Oct. (W-827050) (j-2) – 8 Nov. (PK-663115) (j-1)

▪ **2013** (1.596.896)

TW0 peaks – 30-31 Oct. 1367464) (j-1) – 7 Nov. (PK-94832) (j-2)

▪ **2014** (1.425.747)

ONE wave – 23-24-(25)-26-27 Oct. (W-1040468) (j-1)

▪ **2015** (1.895.072)

TW0 peaks TW0 waves – 20 Oct. (PK-184230) (j-1) - 25-26 Oct. (PK-112473) (J-2) - 30-31 Oct.-1-2 Nov. (W – 1213297) (j-1) – 6-7-8 Nov. (W-222990)

▪ **2016** (1.366.943)






TW0 waves – 19-20-21-22 Oct. (W-459321) (j-1) - 29-30-31 Oct.-1-2 Nov. (W-854195) (j-2)

▪ **2017** (1.217.395)

TW0 waves ONE peak – 23-24-25-26 Oct. (W-551030) (j-1) – 29-30-31 Oct. (W-468816) (j-0) – 7 Nov. (PK-110980) (j-1)

▪ **2018** (2.820.385) *record/20y

TW0 peaks – 23 Oct. (PK- 1073055) (j-1) – 3-4 Nov. (PK – 1638482) (j-1)



Comment

In Pyrenees (France) from 1999 to 2018 (20 years) the

Observers GIFS counted Woodpigeons in transit 31.169.768
(average/y 1.558.488 – min.1.028.000 – max. 2.820.263)

- 816.493 during 52 waves/peaks (average 592.624) – 98,8
{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595
a88d1abe5} of the total 31.169.768
- 227.584 during 21 waves (days more than 2 – average/w
772.242) 52,06
{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595
a88d1abe5} of 31.169.768 (total 20y) and 52,65
{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595
a88d1abe5} of 30.816.493 (total of waves and peaks)
- 588.909 during 31 peaks (1-2 days – average 470.609) –
46,80
{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595
a88d1abe5} of 31.169.768(total 20y) and
47,34{a6be835d09a84c463fa955f66368ad74671de57ffb293aa268
34595a88d1abe5} of 30.816.493 (total of waves and
peaks)

Comparing the averages between numbers of Woodpigeons recorded
during Waves (772.242) and Peaks (470.609) the difference
is great with a
{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1a
be5} increment of
64,1{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a8
8d1abe5} between Peaks and Waves

During first 10 years (1999-2008) Woodpigeons counted
19.517.102 (average/y1.951.710)

During second 10 y (2009-2018) Woodpigeons counted 11.652.660 (average/y 1.165.266)

during the second ten years (2009-2018) the decrement has been 67,5{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5}

Comparing the two decades ,the increment in Falsterbo (Sweden origin/breeding area) 78,8{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} as indicative of growing up of the Scandinavian populations , and the decrement in Pyrenees (transit area to Extremadura-Spain-Portugal) 67,5 {a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} ,it seems that it's possible interpreting this difference : despite the growing up of the Species at the origin areas and at monitoring sites on Mediterranean flyway , most populations prefer to stop wintering in France reducing the length of migration . Anyhow the record of 2018 (2.820.325 in two peaks)seems contradicting this ten-years-trend .

GRAPHICS (2014) at
<https://plus.google.com/photos/103942035281038458760/albums/6086432829554328737>

Concerning the specific focus on Air Pressure changes as stimulus to depart (decision making to takeoff by peaks/waves) for Migration (timing from origin and stop-over sites) –apart all the data related to all abiotic and biotic factors as many reported in graphics- we can underline some conclusions previously reported (...Cavina) and can be assimilated to the current updating (2014-2018).

Commenting on the reading table – graphics (2014) of the results, we highlight some aspects –almost specular as for

Falsterbo – more related to abiotic factors, analyzed on the two temporal “48h/36/24h” and “18h”

prior to take-off.

We note:

- Weather conditions almost never ever disturbed and mostly clear-cloudy;

- Winds of moderate strength with rates prevailing directions from the East to Southeast;

- The maximum visibility always 6-10 km;

- Length of day between 9:30 a.m.-11h ‘and dawn hours. 7.30 to 8 am

- Almost always mild temperature

- Humidity greater than 81{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} in about 70 5 peaks/waves

- Phases of the moon with the illuminated surface of the moon more than 40{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} in 56,76{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of peaks/waves.

- ATMOSPHERIC PRESSURE or AIR PRESSURE (AP) : in 92,85{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of the peaks/waves there was increase (1-30 hPa) with variations above 10 hPa in 75.80{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a

88dlabe5} of the peaks/waves for the sector “36/24 h” and 76.19{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88dlabe5} for the sector

“18 h” before the take-off.

All this confirms the analysis-as “Sweden” – that even in the decision to terminate the stop (stop-over) more or less prolonged the increase in atmospheric pressure event is almost absolute and most important of entities, and what constitutes an acquisition of absolutehighlight: the increase of the atmospheric pressure likely identified with “the finger pressing “ result is not absolute because the 7.14{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88dlabe5} of the peaks/waves atmospheric pressure did not increase or even decreased. And possible variables must be considered attributable to the environmental conditions of the parking influential on abiotic and biotic factors. It should be noted therefore further confirms that the atmospheric pressure is the safety factor for the decision to takeoff to migration and realized more specifically in the afternoon / night before take-off: 90.47{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88dlabe5} of the peaks/waves in the atmospheric pressure is higher than 1010 hPa and more sectorally in 50.70{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88dlabe5} (previous day) and 66{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88dlabe5} (Day of the take-off) is higher than the atmospheric pressure at 1020 hPa. The data of 66{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88dlabe5} (day of Take.Off) of the stop-over France should be put in comparison with similar 53.43{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88dlabe5} in Sweden. Similarly, the investigation of Falsterbo-Sweden, we extrapolated 15 peaks/waves with magnitude of transit above the global average of 42 peaks: in

this case we could not detect significant differences with respect to the overall characteristics of all the peaks/waves .

Some conclusion-comment about Sweden-France's data : during a long period of 20 years (1999-2018) 44.830.138 woodpigeons (Columba palumbus) were recorded by Observers Researchers of official local Institutions and 112 peaks/waves have been identified by retroactive licensed analysis connected with precedent work (1999-2013 results)(...) and more than 90 {a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of the migration-takeoffs happened after a significant documented growth of Air Pressure 12-48 hours before the events .

ITALY-by Club Italiano del Colombaccio “ Progetto Colombaccio”
(41 – 31)

<https://www.ilcolombaccio.it/CMS/>
<http://journal.ilcolombaccio.it/>

If the above-mentioned materials-methods-results were related to foreign experiences by high levels of Research and Institutional organization , in Italy a similar experience monitoring the woodpigeons' migration started by the single pioneering enterprise of Rinaldo Bucchi (Club Italiano del Colombaccio) with the sharing of more than hundred hunters-observers (“Citizen science” policy) member of Club Italiano del Colombaccio , in a research-project named in 1997 “ Progetto Colombaccio “.

The data collected monitoring the woodpigeons' migration in Italy (Citizen science) for 10 year became basic for the dignity of an International paper (35) :

< **Cavina E.,Bucchi R.,Busse P. – (2018) – The General Pattern of Seasonal Dynamics of the Autumn Migration of the Wood Pigeon (Columba Palumbus) in Italy – THE RING 40 (2018) 10.1515/ring-2018-0001** and

<https://www.researchgate.net/publication/328336987> >

where the analysis of peaks and waves has defined a standard methodology on the specific item. The same data have been previously reported to be compared with the Sweden-France results by focus on Air Pressure changes and migration-peaks-waves in Italy .

ITALY data collected in Archives of “Progetto Colombaccio” Club Italiano del Colombaccio (Head of the Team Rinaldo Bucchi) – Official Migration counts on 4.831.138 woodpigeons counted 1998-2007 (10 years) – 12 peaks/waves of migration .
Details of the analysis are in <http://www.scienceheresy.com/ornithologyheresy/Cavina2015.pdf>

ITALY 1998-2007 : 12 peaks/waves

4.831.138 birds (annual average 483 831)

-1998 – 15 October (8 days)

-1999 – 27 October (2 days)- 8 November(4 days)

-2000 – 14 October (8 days)

-2001 – 11 October (5 days)

-2002 – 20 October (2 days)

- 2003 – 16 October (2 days)
- 2004 – 19 October (2 days)
- 2005 – 12 October (6 days)
- 2006 – 26 October (3 days)- 4 November (3 days)
- 2007 – 15 October (2 days)

In November we have considered only two significant peaks over 10 .

Comment

About the data collected in Italy we can detect important elements as following.

In the period 1998-2007 (ten years) with the collaboration of 115 observers of the Club of Woodpigeon, were surveyed in transit 4.831.138 birds (annual average 483 831) and have been identified 10 migratory PEAKS/waves (min.1g. -max.6gg) in October and 10 peaks/waves in November (here used two). Hours of transit were: rise - 9 am (52{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5}), 9 - 14 h (25{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5}), 14 - h night (23{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5}) Consistency of flights observed: from 1 to 50 birds (82{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5}), from 51 to more than 300 (18{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5}) Average number of annual observation point: 3,963 birds Age (only from game bag):

20{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} young,
 28{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of young adults
 52{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} The “waves of step” (fury) account for
 15-20{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5} of total migration (as questionable and
 approximate). The maximum value (annual) of sightings was in
 Veneto and Friuli (door Migration Center-East) in 2002:
 18,683birds ; and the minimum value in the Basilicata region
 in 2003: 416 birds.

Concerning the Air Pressure we were confronting FOUR AREAS
 along the MIGRATION ROUTE from “nesting areas “ and “stop-over
 areas” (possibly mixed) on Long – Medium – Short distance (A-B-C) arriving to the “transit area in Italy” (D), with the
 following results about

- Changes in atmospheric pressure (AP) 48/36/24 h prior to
 takeoff in the areas of origin
- VALUES absolute core of the PA in the 12 h prior to takeoff
 in those areas.
- CHANGES: -in the long distance remote areas (more than 1500
 km) he A.P. increased (78-48 h

before transit in Italy) 67
 {a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1abe5};

-in the most remote and continental stop-over areas (Medium
 and short distance) in the A.P.

48/36/24 h prior totakeoff and the day of detection of the
 peak (in ITALIAN) rises by more than 10

hPa in 90
{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1a
be5} of the Peaks/waves

– The same: in the intermediate areas and closer to the coast
(short distance – 24-12 h. before

take.off) , the most significant of " takeoff of the day
“(latitudes 47 ° 30 ‘, 46 ° 03’, 45 ° 20 ‘, 41 °

53 ‘), the increase is more than
90{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d
1abe5}

-Values Absolute: in the most remote and interior the basic
values of PA – 12 hours before – are

higher than 1010 hPa in the
75-100{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595
a88d1abe5}

– The same: in the intermediate areas the basic values greater
than 1010 hPa are in the
50{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d
1abe5}

-66{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88
d1abe5}

and in the areas closest coastal
83.33{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a
88d1abe5}

– Check-in (ITALY) the absolute values of the basis on the
first day of the peak/wave are higher than 1010 hPa 50
{a6be835d09a84c463fa955f66368ad74671de57ffb293aa26834595a88d1a
be5}.

All of these data-elements – found, as mentioned above, with
precise detail (day by day, hour by hour) taken from the
website Weather History (historical archives) freely available

on the Internet, and transferred to paper ballots useful for comparative examination – there seem to indicate clearly that trust with the abiotic factor most decisive fledging mass migration (fury) is readily identifiable in the “overhang” – more or less progressive or sudden, but always such – above 10 hPa 48-12h before takeoff. “ It ‘important that this data “ITALY” (4.83138 million census – 1998 to 2007 – 12 Peaks/waves) – particularly for 48-36 h before the start of raising – line up with a similar finding in the Pyrenees (census 27,354,388 Pigeons 1999-2013 – 42 peaks) being the areas of takeoff of the first day of peak/wave, all areas of prevailing stop-over more or less prolonged. More properly in the Nordic nesting area (Sweden) (10,760,141 census Pigeons -1999-2013 – 47 peaks) the “overhang” of PA is steeper and closer to the time of the first takeoff mass migration (12- 24 h before). It should be noted that all data collected are drawn from a global basis (Sweden-Italy-Pyrénées) of 42,945,909 woodpigeons actually surveyed in transit flight in 10-15 years and featuring 101 occasions flying of peak migration (1- 6 days).

The ” **temporal extension of the Research** “(November 2018) has enabled an analysis of both the

predictions of both the actual feedback directly in the field (“ **on the field** “) in the more recent migratory seasons, realizing itself as an analysis of” live “, that can also be critical when viewed in the rigidframework of a scientific research, but in fact collect tangible proofs of the relationship between abiotic factors (including atmospheric pressure) and massive takeoff and mass flows in the migration.

In 2014 in ITALY: the performance of the full migration (with “peaks”) was characterized by differences in weather

conditions between the Tyrrhenian Sea area (cyclonic vortex beginning of October) and the Adriatic Sea area (entrance from the Balkans – High Pressure permanent) with – Early mass influx of October 7 to 9 on the side of the Adriatic Sea – More late influx of mass 23 to 24 October on the Tyrrhenian side Very important were the massive takeoffs observed directly (<http://www.ilcolombaccio.it/>): – 19- October 23 from the woods of St. Rossore (Pisa-Livorno) * – October 20 forests in Adriatic area -PesaroUrbino – October 25, Mesóla forest (Ferrara) * – 25 to 27 October Monte Conero (Ancona) – November 2 (flow) – Grosseto area Tirreno The phenomena of takeoff “strip” of thousands and thousands of pigeons, at dawn, and S.Rossore and Mesola have been documented with videos and photographs. All mass migration (2014) – as regards the abiotic factors – were characterized by gradual growth conditions (stability of high atmospheric pressure equal to or greater than 1020 hPa) or rapid increase of 36-12 h. prior to fly both on the Adriatic Sea both on the Tyrrhenian Sea All mass takeoffs migration from the Italian stop-over areas were characterized by rapid increases in PA with changes (24-12 h) of more than 10 hPa.

Many other documented takeoffs (2014-2018) after jumps of Air Pressure more than 10 hPa in the stop-over area in Italy are documented in <https://www.ilcolombaccio.it/CMS/category/colombaccio-scientifico/>

Many of these takeoffs have been forecast by the daily control of forecast Air Pressure graphics in many Meteo-websites : the precision of forecasts is an important support to the scientific methodology .

It's emblematic the takeoff of 200-300.000 woodpigeons 31 October 2017 from Mesola forest (Ferrara-Italy) fully documented and by the video <https://youtu.be/pGK6z9SY8Cg> (Denis Bianchi -2017) and our forecast 4 days before the event

In 2017 the Club Italiano del Colombaccio has produced an experimental project by a WebApp (MCL) to monitoring daily/hourly the autumn migration by cooperation of registered Observers on a Italy'map real-time on-line . The MCL results of 2017 (37) are at <http://www.scienceheresy.com/ornithologyheresy/Woodpigeons.pdf>

7 -9 Oct. – One wave – 40.795 woodpigeons by 676 sightings –AP jump 13 hPa

25-26 Oct–One peak – 55.945 woodpigeons by 645 sightings – AP jump 23 hPa

31-Oct.-! Nov.-One peak – 205.950 woodpigeons by 172 sightings- AP jump 27 hPa



In 2018 MCL as experimental method has encountered some technical problems ,but some data are indicative

11-15 Oct. – One wave – 58.615 woodpigeons – AP jump 15 hPa

20-21 Oct. – One peak – 37.880 woodpigeons – AP jump 21 hPa.

All (2017-2018) peaks/waves have been previously predicted by AP-meteo-forecast and focus on AP forecast-graphics 5 days

before .

All the ITALY's data (1999-2007 and 2014-2018) – also directly from the field as in the last years – confirm the prevalent influence of AP jumps on the migration's "decision making" .

Other retroactive analysis of relationships between AP changes and birds' migration has been reported for other 4 birds' specie as by "THE PARA-TYMPANIC ORGAN of VITALI: the challenge of bird sensory physiology." 2016 – <http://www.scienceheresy.com/ornithologyheresy/Cavina2016.pdf> (39)

"we analyzed abiotic factors on the origin Scandinavian-Russian breeding areas for four more bird species in addition to the Woodpigeon. We have selected 60 migratory mass peaks of transit of the Eurasian Siskin, the Chaffinch/Brambling, the Common Starling, the Common Woodpigeon and the Barnacle Goose. The detailed relationship between mass take-off and changes in Atmospheric Pressure (plus other abiotic factors) 48-12 hours before starting has been analyzed . – (A) – 6 peaks not significant – (B) – 6 peaks moderately significant (7-10 hPa difference) – (C) – 48 peaks strongly significant (10 – 20 hPa difference) The supposed role of the Para-Tympanic Organ of Vitali (PTO) is underlined and further investigated."

documented "evidence based" data concerning the Text are available at

https://photos.google.com/share/AF1QipM79yvIsGsL0iiwU_wuU3evW2-Kpqk1jTfMZwQ6dw6zRUS44mmb0S2dA2uCXPn5Ag?key=QXFxeDMzVUV0WTcxbnNlRG43RlJhYjJ0cXlRbnh3

DISCUSSION

Some elements of above-mentioned items (comments) have been already directly discussed. (41-39)

Taking attention to the target of the present paper (PT0 and scientific investigations) other items must be considered .

- PT0 has been discovered more than a century ago by a single genial Researcher using no more than a microscope and histological procedure of that time .(1)
- Identification of specific supposed physiological relationships was genial already (barometric/altimetical sense).(1-28)
- Controversial items have been developed at that time but the specific barometric supposed function remained valid . (14-15)
- Many specific research fields are unexplored and still not investigated.
- Evolution's implications from fishes to birds are well documented by many researches and are still fascinating.(30)

Concerning this last item ,many evolutionary theories about sense-organs have considered PT0.

“ The amniotic PT0 develops from a previously undiscovered sensory placode “(O’Neill 2012....)

“PT0 stems from a unique placodeplacodes are specialized patches of thickened embryonic skin from wich sense organs developit will be interesting to use molecular techniques to determine whether the PT0 placode starts to develop in mammals , even if this organ itself is not formed” “ After initial excitement wich led to Vitali being nominated for the

1934 Nobel Prize in Physiology or Medicine , its existence was largely forgotten and ignored “(Baker C. – Sept.2012...) and we can affirm “ still in present 2018 time forgotten and ignored to be able to attract curiosity and research in modern sense's ecology “ .

Other generalist consideration is in Web-literature : “ During the middle part of 20th Century the strange story of the <organ of flight> passed quietly into the annals of Science and was almost forgotten” . New technologies show the ability of birds to move themselves in changing Air Pressure conditions “ but something else is needed. Remember Prof.Vitali's PT0 ? Let take a closer look at it.... And –although conclusive proof still waits on some ongoing research – many Scientists are becoming convinced that this (PT0) is the missing piece of the puzzle . Satellite tracking has shown that **long-distance migrants can maintain a constant altitude**they are able to do this within a few tens of meters ,also in total darkness ...the only way this could be possible is if birds have a built-in “altimeter” and PT0 as Air Pressure sensor fits the bill perfectly “ ...and “ sensing the weather in advance” (by Skyscanner Paul Wildlifewriter – 14 Sept.2013 < The weather and
migrating birds
–<http://thewildlifewriter.blogspot.com/2013/09/skyscanner.html>
>”

The specific Literature'references' support remains poor after (2014-2018) full Bibliography overview reported by Giannessi F. (8) and our updating (41) .

At the same time it seems unacceptable to us that relatively ancient (1950) and almost modern (1998) papers have not considered PT0 in discussion and bibliography . Most recent sample of this forgotten bibliography is an excellent paper concerning altitude and fly of birds at twilight exploring in depth the relationships with AP changes in altitude and no mention about PT0 (Behavioral Ecology and Sociobiology

(2018)72.45 – Swiss Ornithol. Inst.) : it could be right remember the motto of Prof. Vitali “ no science without anatomy “.

Apart the basic items discussed and related comments concerning the status of knowledge about PT0 physiology , we have reported updated details in a precedent paper on-line as *“Finally, we emphasize that the sensor terminal of the changes in atmospheric pressure can be easily detected in the organ Para-Tympanic (PT0) Vitali, which studied for the first time by Vitali in Italy in the early decades of the last century, still the subject of extensive research morphological and functional :if “the finger pressing” can be discerned in the changes of atmospheric pressure (the “overhang”), “click” on which the press is probably the Paratympanic organ of Vitali, having to consider all the neuro-functional integration with the adjacent structures in the inner ear (Lagena, vestibular apparatus) until the centers of the Brain and Cerebellum “*

Anatomy

:

<https://plus.google.com/photos/103942035281038458760/albums/5802521945641185121>


von Bartheld CS.- Functional morphology of the paratympanic organ in the middle ear of birds. –Brain Behav Evol. 1994;44(2):61-73.-

<http://www.ncbi.nlm.nih.gov/pubmed/7953609>

By this way the discussion can evolve to anatomo-physiologic items .

ANATOMY at the special Link

<https://plus.google.com/photos/103942035281038458760/albums/5802521945641185121>



In a current analysis and bibliographic review concerning anatomo-physiologic elements of PTO , we are surprised to discover that nothing is described or studied about the fluid (muco-gelatinous) present inside PTO saccula wich obviously has a role to transmit AP changes and forces (including gravitational,electromagnetic,moon phases variations) to the sensitive hair-cells of PTO. These molecular elements of this fluid (as +/- iones,Ca,K,proteins'architecture) will be very probably similar to the electro-chemical composition of the fluids in the inner ear (*Thalmann I, Hughes I, Tong BD, Ornitz DM, Thalmann R. Microscale analysis of proteins in inner ear*

tissues and fluids with emphasis on endolymphatic sac, otoconia, and organ of Corti. Electrophoresis. 2006;27(8):1598-608.) . These elements and their electrochemical functional activity could be specific to “inform” the hair-cells sensors about the AP changes ,together the physical force received and transmitted by the tympanic membrane and its ligaments connected with PT0 .All these electro-chemical-molecular interactive elements are still unexplored .The ancient and present (1986-2010) knowledges concerning the fluid function in inner ear and acustic and vestibular physiology could be a guide-line to develop new researches focused on PT0’s fluid. (ISVR Technical Report 135, 1986 The effect of changes in cerebrospinal fluid pressure on the labyrinth in terms of tympanic membrane displacement. Tweed JJ, Marchbanks RJ and Martin AM. ISVR, Southampton University, Technical Report 135, 1986 -//- Maria del Socorro Hernandez-Montes Study of tympanic membrane displacements with digital holographic interferometry [Proceedings Volume 7387, Speckle 2010: Optical Metrology](#); 73871C . ISVR, Southampton University, Technical Report 135, 1986 -//- Maria del Socorro Hernandez-Montes Study of tympanic membrane displacements with digital holographic interferometry [Proceedings Volume 7387, Speckle 2010: Optical Metrology](#); 73871C)

Remaining on the same electrophysiological field concerning transmission and modulating stimulus ,better ultrastrucutral studies on the structures of PT0 hair-cells and their nerves’ connections – also with Lagen, cochlear system, brain’s area – perhaps could be developed by new electro-microscopical technologies

([https://www.cell.com/current-biology/pdfExtended/S0960-9822\(17\)30730-3](https://www.cell.com/current-biology/pdfExtended/S0960-9822(17)30730-3) Electron Microscopic Reconstruction of Functionally Identified Cells in a Neural Integrator) ..

The complexity of all these questions/answers related to “AP changes &PT0” is evident and open to better modern researches

.

Many other strictly ornithological research items could be discussed as by ornithological research model proposed “on the field “ (experimental behaviour monitoring in wild birds on the field , related to AP changes) (“La Migrazione autunnale del Colombaccio in Italia “ book&E-book by Aracne Editrice – Rome-Italy – 2018) (36)

CONCLUSION

The present paper together with the personal bibliography (2013-2018) could be easily marked as a fruit of a fantasy-work of a dilettante . We try to affirm that this work could be accepted as a work managed by a scientific personal background of the Author supported by the “Citizen science” (Italian Journal Woodpigeon Research – Journal on-line – <http://journal.ilcolombaccio.it/>) spirit of many members of Club Italiano del Colombaccio .

The collecting data methodology and analysis have been developed for many years by serious work on the field and on the writing tables . It was heavy but managed by determination : all the reported data are true and many times controlled and discussed also with criticism.

The “evidence based” data – also if reported by raw numbers – shows the evidence of the decisive role of Air Pressure changes in the “decision making” of birds Migration in Europe and in Italy .

So important stimulus –together with other abiotic-biotic factors – must have a body-target-platform where itself stimulus can put its force .

New technologies(Satellite – radio-tracking) show the ability of birds to move themselves in changing Air Pressure

conditions : therefore in the body of the birds some sensors must be able to receive and manage this physic stimulus .

More than a century ago Giovanni Vitali discovered the Paratympenic Organ as a new sensitive organ linked with barometric and altimetric functions , expression of this physiologic evolutive destiny from fishes (prufondimeter) to birds (altimeter) (28- 30- 20-19)

Despite many sporadic (one Century) and documental researchers' efforts (6 > 34),conclusive proof still waits to be accepted . "Evidence based" data of our retroactive and real-time research in Europe and Italy (wild birds) (35 > 43) should be able to convince new Researchers and Institutions to develop new investigations and new methods to solve in depth the PTO mystery .

The appeal of F.Giannessi (2014) (8) is still unheeded : *"PTO is used daily by enormous number of living animals (200-300 billions of birds) but represents currently one of the major unsolved mysteries and challenges in sensory physiology of vertebrates "* .

The mystery of <https://youtu.be/pGK6z9SY8Cg> is still open to be investigated .

Our position of "dilettante" Author of the above – text , let us free from Academic liturgical obligations : we have not and never will have –as dilettante- the power to develop suitable new institutional specific Research's efforts , but we have the freedom to cry our aware indignation to the Science .



BIBLIOGRAPHY and Web-bibliography

- Vitali G. Di un interessante derivato della prima fessura branchiale nel passero. Anat Anz. 1911;39:219–224.
- Vitali G. Di un interessante derivato dell'ectoderma della prima fessura branchiale nel passero. Un organo nervoso di senso nell'orecchio medio degli uccelli. Anat Anz. 1912;40:631–639
- Vitali G. Il comportamento dell'organo della prima fessura branchiale (placode epibranchiale) nei Selaci. Monit Zool Ital. 1925;36:122–130.
- Vitali G. Sui disturbi funzionali e sulle lesioni

istologiche dipendenti della distruzione dell'organo nervoso di senso da me descritto nell'orecchio medio degli Uccelli. Atti dell'Acc Dei Fisiocritici di Siena. 1915a;5 (7):291–302

- Vitali G. Il comportamento dell'organo della prima fessura branchiale negli anfibi, nei rettili e nei mammiferi. Ricerche Morfol. 1924;4:191–220.

- **Nomination Database. Physiology or Medicine. 1934 6 Dec 2011**http://www.nobelprize.org/nobel_prizes/medicine/nomination/nomination.php?action=show&showid=1990

- Ruffini A. Sull'organo nervoso paratimpanico di G. Vitali od organo del volo degli uccelli. Boll Sci med Soc med-chir Bologna. 1920:141–154.
Archo ital Otol Rhinol Lar. 1920;31:397–413

- [Francesco Giannessi](#),^{a,*} [Riccardo Ruffoli](#),^a and [Christopher S. von Bartheld](#) – (2013) –

Giovanni Vitali: Discoverer of the Paratympanic Organ – [Ann Anat. 2013](#)

Jan;195(1): 10.

- von Bartheld C.S., Giannessi F.- (2011)- The paratympanic organ: a barometer and altimeter in the middle ear of birds?- *Exp. Zool. (Mol. Dev. Evol.)* 316:402–408, 2011
- Giannessi F. On the presence of reciprocal synapses in

the paratympanic organ of the chicken. Anat Embryol (Berl) 1989;180:175–178.

- Giannessi F. Ultrastructure of the supporting cells of the paratympanic organ in the chicken: a preliminary study. Boll Soc Ital Biol Sper. 1990;66:99–104
- Giannessi F, Ruffoli R. The ultrastructure of the sensory hair cells of the paratympanic organ receptor cells in chicken. Anat Embryol. 1996b;193:569–575
- Federici F. Ueber die Innervation des von Vitali entdeckten Sinnesorgans im Mittelohr der Voegel (sogen. paratympanisches Organ) Anat Anz. 1927;62:241–254.
- Benjamins CE. Y a-t-il une relation entre l'organe paratympanique de Vitali et le vol des oiseaux? Arch Neerl Physiol. 1926;11:215–222.
- Benjamins CE. L'organe paratympanique de Vitali, soi-disant organe du vol des oiseaux, chez le pingouin. Acta otolar. 1939;27:266–270.
- Kreithen ML, Keeton WT. Detection of changes in atmospheric pressure by the homing pigeon, (*Columba livia*. J
- Giannessi F, Pera L. On the presence of fibers of probable efferent function in the paratympanic organ in chickens. Boll Soc Ital Biol Sper. 1987;63:337–340
- Giannessi F, Fattori B, Ruffoli R, Gagliardo A. Homing experiments on pigeons subjected to bilateral destruction of the paratympanic organ. J Exp Biol. 1996;199:2035–2039.
- Barry MA, Boord RL. The spiracular organ of sharks and skates: anatomical evidence indicating a mechanoreceptive role. 1984;226:990–992.

20) Baker CVH, O'Neill P, McCole RB. Lateral line, otic and epibranchial placodes:

developmental and evolutionary links? J Exp Zool B Mol Dev

Evol. 2008;310:370–383.

- Jaskoll TF, Maderson PFA. A histological study of the development of the avian middle ear and tympanum. *Anat Rec.* 1978;190:177–200.
- Neeser JA, von Bartheld CS. Comparative anatomy of the paratympanic organ (Vitali organ) in the middle ear of birds and non-avian vertebrates: focus on alligators, parakeets and armadillos. *Brain Behav Evol.* 2002;60:65–79
- Petrash EW, Andres KH, von Düring M, Delius JD. Morphologische Aspekte des Vitali-Organs bei der Taube. *Anat Anz.* 1983;153:288.
- Ruffoli R, Giambelluca MA, Giannessi F. Ultrastructure of the supporting cells in the paratympanic organ of chicken, *Gallus gallus domesticus*. *J Morphol.* 1998;236:65–73.
- Simonetta A. L'organo di senso dello spiracolo e l'organo paratimpanico nella sistematica dei vertebrati. *Arch Ital Anat Embryol.* 1953;58:266–294
- von Bartheld CS. Development and innervation of the paratympanic organ (Vitali organ) in chick embryos. *Brain Behav Evol.* 1990;35:1–15
- von Bartheld CS. Functional morphology of the paratympanic organ in the middle ear of birds. *Brain Behav Evol.* 1994;44:61–73.
- von Bartheld CS, Giannessi F. The paratympanic organ: a barometer and altimeter in the middle ear of birds? *J Exp Zool B Mol Dev Evol.* 2011;316:402–408. [[PMC free article](#)]
- von Bartheld CS, Rubel EW. Paratympanic and spiracular sense organs: phylogenetic distribution and theories of function, including hearing. In: Webster DB, Fay RR, Popper AN, editors. *Evolutionary Biology of*

Hearing. Springer; New York: 1992. pp. 582–583.

- O'Neill P. Molecular mechanisms underlying paratympanic organ development and evolution. RIKEN FY2009 Foreign Postdoctoral Researchers. 2010;22:193–194.

31) Bucchi R. - (2018) – **Selective Migration Monitoring (MSM) of Woodpigeons migrated in**

Italy

<http://journal.ilcolombaccio.it/ijwr-vol-1-2018-papers-short-communications/>

32) Martin G. – (2017) - The sensory ecology of birds – Oxford Univ. Press 2017

33) Alerstam T. - (1997) - Bird Migration – Cambridge Univ. Press 1997

34) Newton I. - (2007) – The migration ecology of birds – Elsevier Pub. 2007

CAVINA E. – bibliography ***

35) Cavina E., Bucchi R., Busse P. – (2018) – The General

Pattern of Seasonal Dynamics of the Autumn Migration of the Wood Pigeon (Columba Palumbus) in Italy – THE RING 40 (2018) 10.1515/ring-2018-0001and

<https://www.researchgate.net/publication/328336987> *****

36) Cavina E.,Bucchi R.,Bianchi D.,Giovanetti G.,Feligetti V.,Giannerini S.,Bececco L –

La Migrazione autunnale del Colombaccio (*Columba palumbus*) in Italia – Ed.

Aracne Editrice-Roma – 2017

37) Enrico Cavina, Denis Bianchi, Vasco Feligetti, Graziano Giovanetti and Rinaldo Bucchi- (2017)- Monitoring the 2017 Autumn Migration of the Woodpigeon (*Columba palumbus*) : Take-off decision making and forecasting .
<http://www.scienceheresy.com/ornithologyheresy/Woodpigeons.pdf>

<http://journal.ilcolombaccio.it/ijwr-vol-1-2018-papers-short-communications/>

38) Cavina E. – (2017)- The Para-Tympanic organ of Vitali : Research's failure or

challenge ?

<http://www.scienceheresy.com/ornithologyheresy/VitaliPT0.pdf>

39) Cavina E. –(2016)- THE PARA-TYMPANIC ORGAN of VITALI: the challenge of bird

sensory physiology .

<http://www.scienceheresy.com/ornithologyheresy/Cavina2016.pdf>

41) Cavina E. –(2014) Decision making of autumn migrations of woodpigeons (*Columba*

palumbus) in Europe: analysis of the abiotic factors and atmospheric pressure changes

<http://www.scienceheresy.com/ornithologyheresy/Cavina2015.pdf>

42) Cavina E. , Cenni P. –
Woodpigeons' (*Columba palumbus*) autumn migration in

Italy monitored
(consecutive 19 years) in a single
crucial spot by single Observer and

uniform method.
– *Preliminary report*

<http://journal.ilcolombaccio.it/ijwr-vol-1-2018-papers-short-comunications/>

43) Cavina E.,Bucchi R. – (2018) – Woodpigeons' (*Columba palumbus*) autumn in eight

migration in Central and Northern Italy along two flyways monitored (*consecutive 7*

years) crucial spots by eight Observers and uniform method .

<http://journal.ilcolombaccio.it/ijwr-vol-1-2018-papers-short-communications/>

45) Anatomy of Woodpigeons and special focus on PT0 – Photographic Atlas of Enrico Cavina

<https://plus.google.com/photos/103942035281038458760/albums/5802521945641185121>

FOR ANY PROBLEM ON THE WEB-LINKS , please, COPY and PASTE on the window of GOOGLE search motor .